

NUZYRA® ORAL DOSING IN ABSSSI

ONCE-DAILY DOSING¹: Treatment duration: 7-14 days¹

DAY 1-2: Loading Dose

450 mg (3 tablets)
once a day on Days 1 and 2



Tablets are not shown at actual size.

Once-Daily Maintenance Dose

300 mg (2 tablets)



HOW TO PRESCRIBE



NUZYRA 450 mg

PO QD x 2 days

THEN

NUZYRA 300 mg

PO QD

7-14 days total

**TOTAL
TABLETS:
16-30**

THE CURRENT SIS GUIDELINES FOR THE MANAGEMENT OF SSTIs²

Recommend NUZYRA as an alternative treatment for suspected or confirmed MRSA in certain ABSSSIs

Choice of alternative antibiotic should be driven by multiple factors, including clinician experience, history of drug allergies or adverse events, drug interactions, and drug-disease interactions.

Criteria for selecting antibiotic therapy²:

- When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy
- In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy

MRSA=methicillin-resistant *Staphylococcus aureus*; PO=per os; QD=once a day; SIS=Surgical Infection Society; SSTIs=skin and soft tissue infections.

NUZYRA is available today at our network specialty pharmacies

Visit nuzyra.com/find

INDICATION and USAGE


INDICATION: NUZYRA® (omadacycline) is a tetracycline-class antibacterial indicated for the treatment of Acute Bacterial Skin and Skin Structure Infections (ABSSSI) in adults caused by certain susceptible microorganisms. See complete [Indication and Usage](#).

USAGE: To reduce the development of drug-resistant bacteria and maintain the effectiveness of NUZYRA and other antibacterial drugs, NUZYRA should be used only to treat or prevent infections that are proven or strongly suspected to be caused by susceptible bacteria.

IMPORTANT SAFETY INFORMATION

CONTRAINDICATIONS: NUZYRA is contraindicated in patients with known hypersensitivity to omadacycline or tetracycline-class antibacterial drugs, or to any of the excipients.

Please see [Important Safety Information](#) and full [Prescribing Information](#) at [NUZYRA.com](https://nuzyra.com).



once-daily

NUZYRA®
(omadacycline)

100 mg for injection / 150 mg tablets

WHEN PRESCRIBING ORAL NUZYRA[®], INSTRUCT PATIENTS TO¹:



- Fast for at least 4 hours and then take with water
 - NUZYRA can be taken at bedtime or upon waking



- Not eat or drink (except water) for 2 hours after dosing



- Not consume dairy products, antacids, or multivitamins for 4 hours after dosing

Patients on anticoagulant therapy may require downward adjustment of their anticoagulant dosage while also taking NUZYRA.

**PATIENTS WITH RENAL OR HEPATIC IMPAIRMENT
DO NOT REQUIRE A DOSE ADJUSTMENT.¹**

NUZYRA is also available in an intravenous (IV) formulation¹

INDICATION and USAGE

INDICATION: NUZYRA® (omadacycline) is a tetracycline-class antibacterial indicated for the treatment of Acute Bacterial Skin and Skin Structure Infections (ABSSSI) in adults caused by the following susceptible microorganisms: *Staphylococcus aureus* (methicillin-susceptible and -resistant isolates), *Staphylococcus lugdunensis*, *Streptococcus pyogenes*, *Streptococcus anginosus* grp. (includes *S. anginosus*, *S. intermedius*, and *S. constellatus*), *Enterococcus faecalis*, *Enterobacter cloacae*, and *Klebsiella pneumoniae*.

USAGE: To reduce the development of drug-resistant bacteria and maintain the effectiveness of NUZYRA and other antibacterial drugs, NUZYRA should be used only to treat or prevent infections that are proven or strongly suspected to be caused by susceptible bacteria.

IMPORTANT SAFETY INFORMATION

CONTRAINDICATIONS: NUZYRA is contraindicated in patients with known hypersensitivity to omadacycline or tetracycline-class antibacterial drugs, or to any of the excipients.

WARNINGS AND PRECAUTIONS

The use of NUZYRA during tooth development (last half of pregnancy, infancy and childhood to the age of 8 years) may cause permanent discoloration of the teeth (yellow-gray-brown) and enamel hypoplasia.

The use of NUZYRA during the second and third trimester of pregnancy, infancy and childhood up to the age of 8 years may cause reversible inhibition of bone growth.

Hypersensitivity reactions have been reported with NUZYRA. Life-threatening hypersensitivity (anaphylactic) reactions have been reported with other tetracycline-class antibacterial drugs. NUZYRA is structurally similar to other tetracycline-class antibacterial drugs and is contraindicated in patients with known hypersensitivity to tetracycline-class antibacterial drugs. Discontinue NUZYRA if an allergic reaction occurs.

Clostridioides difficile associated diarrhea (CDAD) has been reported with use of nearly all antibacterial agents and may range in severity from mild diarrhea to fatal colitis. Evaluate if diarrhea occurs.

NUZYRA is structurally similar to tetracycline-class antibacterial drugs and may have similar adverse reactions. Adverse reactions, including photosensitivity, fixed drug eruption, pseudotumor cerebri, and anti-anabolic action (which has led to increased BUN, azotemia, acidosis, hyperphosphatemia, pancreatitis, and abnormal liver function tests), have been reported for other tetracycline-class antibacterial drugs, and may occur with NUZYRA. Discontinue NUZYRA if any of these adverse reactions are suspected.

Prescribing NUZYRA in the absence of a proven or strongly suspected bacterial infection is unlikely to provide benefit to the patient and increases the risk of the development of drug-resistant bacteria.

ADVERSE REACTIONS

The most common adverse reactions (incidence $\geq 2\%$) are nausea, vomiting, infusion site reactions, alanine aminotransferase increased, aspartate aminotransferase increased, gamma-glutamyl transferase increased, hypertension, headache, diarrhea, insomnia, and constipation.

DRUG INTERACTIONS

Patients who are on anticoagulant therapy may require downward adjustment of their anticoagulant dosage while taking NUZYRA.

Absorption of tetracyclines, including NUZYRA is impaired by antacids containing aluminum, calcium, or magnesium, bismuth subsalicylate and iron containing preparations.

USE IN SPECIFIC POPULATIONS

Lactation: Breastfeeding is not recommended during treatment with NUZYRA.

References: 1. NUZYRA [Prescribing Information]. Paratek Pharmaceuticals, Inc. 2. Duane TM, Huston JM, Collom M, et al. Surgical Infection Society 2020 Updated Guidelines on the management of complicated skin and soft tissue infections. *Surg Infect (Larchmt)*. 2021;22(4):383-399.



© 2025 Paratek Pharmaceuticals, Inc. All rights reserved.
PARATEK® and the hexagon logo are registered trademarks of Paratek Pharmaceuticals, Inc.
NUZYRA® and its design logo are registered trademarks of Paratek Pharmaceuticals, Inc.
US-NUA-0648-02 04/25

Please see full [Prescribing Information](#) at [NUZYRA.com](#).



once-daily
NUZYRA®
(omadacycline) 100 mg for injection
150 mg tablets